

INFORMAL COMMUNICATION  
Patent Application Docket No. SPO.124

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Examiner : Zachary C. Howard  
Art Unit : 1646  
Applicants : Tadao Saito *et al.*  
Serial No. : 10/561,022  
Filed : December 16, 2005  
Confirm. No. : 5950  
For : Use of Toll-Like Receptor-Expressing Cells

Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

INFORMAL COMMUNICATION

Dear Examiner Howard:

In preparation for our telephone interview tomorrow at 11:00, following are amendments that I would like to propose to claims 2 and 5. I do not remember, should I call you or will you call me?

Proposed claims

2 (currently amended). ~~The method, according to claim 1, wherein said method is used to screen A method of screening for a sample that activates the intestinal tract immune system, wherein the method comprises comprising the steps of:~~

~~(a) assessing whether a plurality of test samples activate the intestinal tract immune system by the assessment method of claim 1; and contacting a test sample with an isolated cell expressing a naturally-occurring mammalian intestinal tract tissue-expressed Toll-like receptor;~~

~~(b) measuring activity of the Toll-like receptor using signal transduction in the cell as an indicator; and~~

(b)(c) selecting from the plurality of test samples those assessed to activate the intestinal tract immune system the test sample as a sample that activates the intestinal tract immune system if the activity of the Toll-like receptor is increased as compared to activity of the Toll-like receptor in a cell not contacted with the test sample.

5 (currently amended). The method, according to claim 4, wherein said method is used to screen A method of screening for microorganisms that activate the intestinal tract immune system, comprising the steps of:

(a) assessing whether a plurality of test microorganisms activate the intestinal tract immune system by the assessment method of claim 4; and preparing an extract from a test microorganism;

(b) contacting the extract with an isolated cell expressing a naturally-occurring mammalian intestinal tract tissue-expressed Toll-like receptor;

(c) measuring activity of the Toll-like receptor using signal transduction in the cell as an indicator; and

(d) selecting the test microorganism as a microorganism that activates the intestinal tract immune system if the activity of the Toll-like receptor is increased as compared to activity of the Toll-like receptor in a cell not contacted with the extract

(e) selecting from the plurality of test microorganisms one or more microorganisms that are assessed to activate the intestinal tract immune system.